

ATTORNEY DOCKET NO. 14014.0323U3

EXPRESS MAIL NO. EL403200449US

PATENT

5

10

15

20

TO ALL WHOM IT MAY CONCERN:

Be it known that we, JOHN CHIORINI, ROBERT M. KOTIN and BRIAN
25 SAFER, citizens of the United States of America, residing, respectively, at 2604 Loma
Street, Silver Spring, MD 20902, 707 Gormley, Rockville, MD 20850 and 1610 Tifton
Dr., Silver Springs, MD 20902, U.S.A., have invented new and useful improvements in

AAV5 VECTOR AND USES THEREOF

30

for which the following is a specification.

09747789-112100
PATENT 68447460

AAV5 VECTOR AND USES THEREOF

This application is a continuation of international application PCT/US99/11958 filed on May 28, 1999, which claims priority to U.S. provisional application Serial No. 60/087029 filed on May 28, 1998. The PCT/US99/11958 international application and the 60/087029 provisional patent application are herein incorporated by this reference in their entirety.

BACKGROUND OF THE INVENTION

Field of the Invention

The present invention provides adeno-associated virus 5 (AAV5) and vectors derived therefrom. Thus, the present invention relates to AAV5 vectors for and methods of delivering nucleic acids to cells of subjects.

Background Art

Adeno associated virus (AAV) is a small nonpathogenic virus of the parvoviridae family (for review see 28). AAV is distinct from the other members of this family by its dependence upon a helper virus for replication. In the absence of a helper virus, AAV has been shown to integrate in a locus specific manner into the q arm of chromosome 19 (21). The approximately 5 kb genome of AAV consists of one segment of single stranded DNA of either plus or minus polarity. Physically, the parvovirus virion is non-enveloped and its icosohedral capsid is approximately 20-25 nm in diameter.

To date 8 serologically distinct AAVs have been identified and 6 have been isolated from humans or primates and are referred to as AAV types 1-6 (1). The most extensively studied of these isolates is AAV type 2 (AAV2). The genome of AAV2 is 4680 nucleotides in length and contains two open reading frames (ORFs), the right ORF and the left ORF. The left ORF encodes the non-structural Rep proteins, Rep40,

Rep52, Rep68 and Rep78, which are involved in regulation of replication and transcription in addition to the production of single-stranded progeny genomes (5-8, 11, 12, 15, 17, 19, 21-23, 25, 34, 37-40). Furthermore, two of the Rep proteins have been associated with the preferential integration of AAV genomes into a region of the q arm of human chromosome 19. Rep68/78 have also been shown to possess NTP binding activity as well as DNA and RNA helicase activities. The Rep proteins possess a nuclear localization signal as well as several potential phosphorylation sites. Mutation of one of these kinase sites resulted in a loss of replication activity.

10 The ends of the genome are short inverted terminal repeats which have the potential to fold into T-shaped hairpin structures that serve as the origin of viral DNA replication. Within the ITR region two elements have been described which are central to the function of the ITR, a GAGC repeat motif and the terminal resolution site (TRS). The repeat motif has been shown to bind Rep when the ITR is in either a linear or
15 hairpin conformation (7, 8, 26).

 This binding serves to position Rep68/78 for cleavage at the TRS which occurs in a site- and strand-specific manner. In addition to their role in replication, these two elements appear to be central to viral integration. Contained within the chromosome 19
20 integration locus is a Rep binding site with an adjacent TRS. These elements have been shown to be functional and necessary for locus specific integration.

 The AAV2 virion is a non-enveloped, icosohedral particle approximately 20-25 nm in diameter. The capsid is composed of three related proteins referred to as VP1,2
25 and 3 which are encoded by the right ORF. These proteins are found in a ratio of 1:1:10 respectively. The capsid proteins differ from each other by the use of alternative splicing and an unusual start codon. Deletion analysis of has shown that removal or alteration of AAV2 VP1 which is translated from an alternatively spliced message results in a reduced yield of infections particles (15, 16, 38). Mutations within the VP3

09717789.112100

coding region result in the failure to produce any single-stranded progeny DNA or infectious particles (15, 16, 38).

The following features of the characterized AAVs have made them attractive
5 vectors for gene transfer (16). AAV vectors have been shown *in vitro* to stably
integrate into the cellular genome; possess a broad host range; transduce both dividing
and non dividing cells *in vitro* and *in vivo* (13, 20, 30, 32) and maintain high levels of
expression of the transduced genes (41). Viral particles are heat stable, resistant to
solvents, detergents, changes in pH, temperature, and can be concentrated on CsCl
10 gradients (1,2). Integration of AAV provirus is not associated with any long term
negative effects on cell growth or differentiation (3,42). The ITRs have been shown to
be the only cis elements required for replication, packaging and integration (35) and
may contain some promoter activities (14).

15 AAV2 was originally thought to infect primate and non-primate cell types
provided the appropriate helper virus was present. However, the inability of AAV2 to
infect certain cell types is now known to be due to the particular cellular tropism
exhibited by the AAV2 virus. Recent work has shown that some cell lines are
transduced very poorly by AAV2 (30). Binding studies have indicated that heparin
20 sulfate proteoglycans are necessary for high efficiency transduction with AAV2.
AAV5 is a unique member of the parvovirus family. The present DNA hybridization
data indicate a low level of homology with the published AAV1-4 sequences (31). The
present invention shows that, unlike AAV2, AAV5 transduction is not effected by
heparin as AAV2 is and therefore will not be restricted to the same cell types as AAV2.

25
The present invention provides a vector comprising the AAV5 virus or a vector
comprising subparts of the virus, as well as AAV5 viral particles. While AAV5 is
similar to AAV2, the two viruses are found herein to be physically and genetically
distinct. These differences endow AAV5 with some unique properties and advantages
30 which better suit it as a vector for gene therapy. For example, one of the limiting

09717789-112100

features of using AAV2 as a vector for gene therapy is production of large amounts of virus. Using standard production techniques, AAV5 is produced at a 10-50 fold higher level compared to AAV2. Because of its unique TRS site and rep proteins, AAV5 should also have a distinct integration locus compared to AAV2.

5

Furthermore, as shown herein, AAV5 capsid protein, again surprisingly, is distinct from AAV2 capsid protein and exhibits different tissue tropism, thus making AAV5 capsid-containing particles suitable for transducing cell types for which AAV2 is unsuited or less well-suited. AAV2 and AAV5 have been shown to be serologically distinct and thus, in a gene therapy application, AAV5, and AAV5-derived vectors, would allow for transduction of a patient who already possess neutralizing antibodies to AAV2 either as a result of natural immunological defense or from prior exposure to AAV2 vectors. Another advantage of AAV5 is that AAV5 cannot be rescued by other serotypes. Only AAV5 can rescue the integrated AAV5 genome and effect replication, thus avoiding unintended replication of AAV5 caused by other AAV serotypes. Thus, the present invention, by providing these new recombinant vectors and particles based on AAV5 provides a new and highly useful series of vectors.

SUMMARY OF THE INVENTION

20

The present invention provides a nucleic acid vector comprising a pair of adeno-associated virus 5 (AAV5) inverted terminal repeats and a promoter between the inverted terminal repeats.

25

The present invention further provides an AAV5 particle containing a vector comprising a pair of AAV2 inverted terminal repeats.

Additionally, the instant invention provides an isolated nucleic acid comprising the nucleotide sequence set forth in SEQ ID NO:1 (AAV5 genome). Furthermore, the

09747789.112400

present invention provides an isolated nucleic acid consisting essentially of the nucleotide sequence set forth in SEQ ID NO:1 (AAV5 genome).

09717789 "112100
The present invention provides an isolated nucleic acid encoding an AAV5 Rep
5 protein, for example, the nucleic acid as set forth in SEQ ID NO:10. Additionally
provided is an isolated full-length AAV5 Rep protein or a unique fragment thereof.
Additionally provided is an isolated AAV5 Rep 40 protein having the amino acid
sequence set forth in SEQ ID NO:12, or a unique fragment thereof. Additionally
provided is an isolated AAV5 Rep 52 protein having the amino acid sequence set forth
10 in SEQ ID NO:2, or a unique fragment thereof. Additionally provided is an isolated
AAV5 Rep 68 protein, having the amino acid sequence set forth in SEQ ID NO:14 or a
unique fragment thereof. Additionally provided is an isolated AAV5 Rep 78 protein
having the amino acid sequence set forth in SEQ ID NO:3, or a unique fragment
thereof. The sequences for these proteins are provided below in the Sequence Listing
15 and elsewhere in the application where the proteins are described.

The present invention further provides an isolated AAV5 capsid protein, VP1,
having the amino acid sequence set forth in SEQ ID NO:4, or a unique fragment
thereof. Additionally provided is an isolated AAV5 capsid protein, VP2, having the
20 amino acid sequence set forth in SEQ ID NO:5, or a unique fragment thereof. Also
provided is an isolated AAV5 capsid protein, VP3, having the amino acid sequence set
forth in SEQ ID NO:6, or a unique fragment thereof.

The present invention additionally provides an isolated nucleic acid encoding
25 AAV5 capsid protein, for example, the nucleic acid set forth in SEQ ID NO:7, or a
unique fragment thereof.

The present invention further provides an AAV5 particle comprising a capsid
protein consisting essentially of the amino acid sequence set forth in SEQ ID NO:4, or
30 a unique fragment thereof.

Additionally provided by the present invention is an isolated nucleic acid comprising an AAV5 p5 promoter having the nucleic acid sequence set forth in SEQ ID NO:18, or a unique fragment thereof.

5 The instant invention provides a method of screening a cell for infectivity by AAV5 comprising contacting the cell with AAV5 and detecting the presence of AAV5 in the cells.

10 The present invention further provides a method of delivering a nucleic acid to a cell comprising administering to the cell an AAV5 particle containing a vector comprising the nucleic acid inserted between a pair of AAV inverted terminal repeats, thereby delivering the nucleic acid to the cell.

15 The present invention also provides a method of delivering a nucleic acid to a subject comprising administering to a cell from the subject an AAV5 particle comprising the nucleic acid inserted between a pair of AAV inverted terminal repeats, and returning the cell to the subject, thereby delivering the nucleic acid to the subject.

20 The present invention also provides a method of delivering a nucleic acid to a cell in a subject comprising administering to the subject an AAV5 particle comprising the nucleic acid inserted between a pair of AAV inverted terminal repeats, thereby delivering the nucleic acid to a cell in the subject.

25 The instant invention further provides a method of delivering a nucleic acid to a cell in a subject having antibodies to AAV2 comprising administering to the subject an AAV5 particle comprising the nucleic acid, thereby delivering the nucleic acid to a cell in the subject.

BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1 shows Heparin inhibition results. Cos cells were plated in 12 well dishes at 5×10^4 cells per well. Serial dilutions of AAV2 or AAV5 produced and purified as previously described and supplemented with 5×10^5 particles of wt adenovirus were incubated for 1 hr at Rt in the presence of 20 $\mu\text{g/ml}$ heparin (sigma). Following this incubation the virus was added to the cells in 400 μl of media for 1 hr after which the media was removed, the cells rinsed and fresh media added. After 24 hrs the plates were stained for Bgal activity.

10

Figure 2 shows AAV2 and AAV5 vector and helper complementation. Recombinant AAV particles were produced as previously described using a variety of vector and helper plasmids as indicated the bottom of the graph. The vector plasmids contained the Bgal gene with and RSV promoter and flanked by either AAV2 ITRs (2ITR) or AAV5 ITRs (5ITR). The helper plasmids tested contained either AAV2 Rep and cap genes (2repcap) AAV5 rep and cap genes with or without an SV40 promoter (5repcapA and 5repcapb respectively) only the AAV2 rep gene (2rep) in varying amounts (1) or (.5) or an empty vector (pUC). The resulting AAV particles were then titered on cos cells. AAV particles were only produced when the same serotype of ITR and Rep were present.

15

20

Figure 3 shows AAV2 and AAV5 tissue tropism. Transduction of a variety of cell types indicated that AAV2 and AAV5 transduce cells with different efficiencies. Equal number of either AAV2 or AAV5 particles were used to transduce a variety of cell types and the number of bgal positive cells is reported.

25

Figure 4 is a sequence comparison of the AAV2 genome and the AAV5 genome.

09717789-112100

Figure 5 is a sequence comparison of the AAV2 VP1 capsid protein and the AAV5 VP1 capsid protein.

Figure 6 is a sequence comparison of the AAV2 rep 78 protein and the AAV5
5 rep 78 protein.

Figure 7 shows the transduction of airway epithelial cells by AAV5. Primary airway epithelial cells were cultured and plated. Cells were transduced with an equivalent number of rAAV2 or rAAV5 particles containing a nuclear localized β -gal transgene with 50 particles of virus/cell (MOI 50) and continued in culture for 10 days. β -gal activity was determined and the relative transduction efficiency compared. AAV5 transduced these cells 50- fold more efficiently than AAV2. This is the first time apical cells or cells exposed to the air have been shown to be infected by a gene therapy agent.

Figure 8 shows transduction of striated muscle by AAV5. Chicken myoblasts were cultured and plated. Cells were allowed to fuse and then transduced with a similar number of particles of rAAV2 or rAAV5 containing a nuclear localized β -gal transgene after 5 days in culture. The cells were stained for β -gal activity and the relative
20 transduction efficiency compared. AAV5 transduced these cells approximately 16 fold more efficiently than AAV2.

Figure 9 shows transduction of rat brain explants by AAV5. Primary neonatal rat brain explants were prepared. After 7 days in culture, cells were transduced with a
25 similar number of particles of rAAV5 containing a nuclear localized β -gal transgene. After 5 days in culture, the cells were stained for β -gal activity. Transduction was detected in a variety of cell types including astrocytes, neuronal cells and glial cells.

Figure 10 shows transduction of human umbilical vein endothelial cells by
30 AAV5. Human umbilical vein endothelial cells were cultured and plated. Cells were

09717789-112100

transduced with rAAV2 or rAAV5 containing a nuclear localized β -gal transgene with 10 particles of virus/ cell (MOI 5) in minimal media then returned to complete media. After 24 hrs in culture, the cells were stained for β -gal activity and the relative transduction efficiency compared. As shown in AAV5 transduced these cell 5-10 fold more efficiently than AAV2.

DETAILED DESCRIPTION OF THE INVENTION

As used in the specification and in the claims, "a" can mean one or more, depending upon the context in which it is used. The terms "having" and "comprising" are used interchangeably herein, and signify open ended meaning.

The present application provides a recombinant adeno-associated virus 5 (AAV5). This virus has one or more of the characteristics described below. The compositions of the present invention do not include wild-type AAV5. The methods of the present invention can use either wild-type AAV5 or recombinant AAV5-based delivery.

The present invention provides novel AAV5 particles, recombinant AAV5 vectors, recombinant AAV5 virions and novel AAV5 nucleic acids and polypeptides. An AAV5 particle is a viral particle comprising an AAV5 capsid protein. A recombinant AAV5 vector is a nucleic acid construct that comprises at least one unique nucleic acid of AAV5. A recombinant AAV5 virion is a particle containing a recombinant AAV5 vector, wherein the particle can be either an AAV5 particle as described herein or a non-AAV5 particle. Alternatively, the recombinant AAV5 virion is an AAV5 particle containing a recombinant vector, wherein the vector can be either an AAV5 vector as described herein or a non-AAV5 vector. These vectors, particles, virions, nucleic acids and polypeptides are described below.

09717789.112100

09717789 "112100

The present invention provides the nucleotide sequence of the AAV5 genome and vectors and particles derived therefrom. Specifically, the present invention provides a nucleic acid vector comprising a pair of AAV5 inverted terminal repeats (ITRs) and a promoter between the inverted terminal repeats. While the rep proteins of AAV2 and AAV5 will bind to either a type 2 ITR or a type 5 ITR, efficient genome replication only occurs when type 2 Rep replicates a type 2 ITR and a type 5 Rep replicates a type 5 ITR. This specificity is the result of a difference in DNA cleavage specificity of the two Reps which is necessary for replication. AAV5 Rep cleaves at CGGT^GTGA (SEQ ID NO: 21) and AAV2 Rep cleaves at CGGT^TGAG (SEQ ID NO: 22) (Chiorini et al., 1999. J. Virol. 73 (5) 4293-4298). Mapping of the AAV5 ITR terminal resolution site (TRS) identified this distinct cleavage site, CGGT^GTGA, which is absent from the ITRs of other AAV serotypes. Therefore, the minimum sequence necessary to distinguish AAV5 from AAV2 is the TRS site where Rep cleaves in order to replicate the virus. Examples of the type 5 ITRs are shown in SEQ ID NO: 19 and SEQ ID NO: 20, AAV5 ITR "flip" and AAV5 "flop", respectively. Minor modifications in an ITR of either orientation are contemplated and are those that will not interfere with the hairpin structure formed by the AAV5 ITR as described herein and known in the art. Furthermore, to be considered within the term "AAV5 ITR" the nucleotide sequence must retain one or more features described herein that distinguish the AAV5 ITR from the ITRs of other serotypes, e.g. it must retain the Rep binding site described herein.

The D- region of the AAV5 ITR (SEQ ID NO: 23), a single stranded region of the ITR, inboard of the TRS site, has been shown to bind a factor which depending on its phosphorylation state correlates with the conversion of the AAV from a single stranded genome to a transcriptionally active form that allows for expression of the viral DNA. This region is conserved between AAV2, 3, 4, and 6 but is divergent in AAV5. The D+ region is the reverse complement of the D- region.

09717789 "112100

The promoter can be any desired promoter, selected by known considerations, such as the level of expression of a nucleic acid functionally linked to the promoter and the cell type in which the vector is to be used. That is, the promoter can be tissue/cell-specific. Promoters can be prokaryotic, eukaryotic, fungal, nuclear, mitochondrial, viral or plant promoters. Promoters can be exogenous or endogenous to the cell type being transduced by the vector. Promoters can include, for example, bacterial promoters, known strong promoters such as SV40 or the inducible metallothionein promoter, or an AAV promoter, such as an AAV p5 promoter. Additionally, chimeric regulatory promoters for targeted gene expression can be utilized. Examples of these regulatory systems, which are known in the art, include the tetracycline based regulatory system which utilizes the tet transactivator protein (tTA), a chimeric protein containing the VP16 activation domain fused to the tet repressor of *Escherichia coli*, the IPTG based regulatory system, the CID based regulatory system, and the Ecdysone based regulatory system (44). Other promoters include promoters derived from actin genes, immunoglobulin genes, cytomegalovirus (CMV), adenovirus, bovine papilloma virus, adenoviral promoters, such as the adenoviral major late promoter, an inducible heat shock promoter, respiratory syncytial virus, Rous sarcoma virus (RSV), etc., specifically, the promoter can be AAV2 p5 promoter or AAV5 p5 promoter. More specifically, the AAV5 p5 promoter can be about same location in SEQ ID NO: 1 as the AAV2 p5 promoter, in the corresponding AAV2 published sequence. Additionally, the p5 promoter may be enhanced by nucleotides 1-130 of SEQ ID NO:1. Furthermore, smaller fragments of p5 promoter that retain promoter activity can readily be determined by standard procedures including, for example, constructing a series of deletions in the p5 promoter, linking the deletion to a reporter gene, and determining whether the reporter gene is expressed, *i.e.*, transcribed and/or translated. The promoter can be the promoter of any of the AAV serotypes, and can be the p19 promoter (SEQ ID NO: 16) or the p40 promoter set forth in the sequence listing as SEQ ID NO: 17.

It should be recognized that any errors in any of the nucleotide sequences disclosed herein can be corrected, for example, by using the hybridization procedure

described below with various probes derived from the described sequences such that the coding sequence can be reisolated and resequenced. Rapid screening for point mutations can also be achieved with the use of polymerase chain reaction-single strand conformation polymorphism (PCR-SSCP) (43). The corresponding amino acid
5 sequence can then be corrected accordingly.

The AAV5-derived vector of the invention can further comprise a heterologous nucleic acid functionally linked to the promoter. By "heterologous nucleic acid" is meant that any heterologous or exogenous nucleic acid, i.e. not normally found in wild-
10 type AAV5 can be inserted into the vector for transfer into a cell, tissue or organism. By "functionally linked" is meant that the promoter can promote expression of the heterologous nucleic acid, as is known in the art, and can include the appropriate orientation of the promoter relative to the heterologous nucleic acid. Furthermore, the heterologous nucleic acid preferably has all appropriate sequences for expression of the
15 nucleic acid. The nucleic acid can include, for example, expression control sequences, such as an enhancer, and necessary information processing sites, such as ribosome binding sites, RNA splice sites, polyadenylation sites, and transcriptional terminator sequences.

20 The heterologous nucleic acid can encode beneficial proteins or polypeptides that replace missing or defective proteins required by the cell or subject into which the vector is transferred or can encode a cytotoxic polypeptide that can be directed, *e.g.*, to cancer cells or other cells whose death would be beneficial to the subject. The heterologous nucleic acid can also encode antisense RNAs that can bind to, and thereby
25 inactivate, mRNAs made by the subject that encode harmful proteins. The heterologous nucleic acid can also encode ribozymes that can effect the sequence-specific inhibition of gene expression by the cleavage of mRNAs. In one embodiment, antisense polynucleotides can be produced from a heterologous expression cassette in an AAV5 vector construct where the expression cassette contains
30 a sequence that promotes cell-type specific expression (Wirak *et al.*, *EMBO* 10:289

09717789 "112100

(1991)). For general methods relating to antisense polynucleotides, see *Antisense RNA and DNA*, D. A. Melton, Ed., Cold Spring Harbor Laboratory, Cold Spring Harbor, NY (1988).

5 Examples of heterologous nucleic acids which can be administered to a cell or subject as part of the present AAV5 vector can include, but are not limited to the following: nucleic acids encoding secretory and nonsecretory proteins, nucleic acids encoding therapeutic agents, such as tumor necrosis factors (TNF), such as TNF- α ; interferons, such as interferon- α , interferon- β , and interferon- γ ; interleukins, such as
10 IL-1, IL-1 β , and ILs -2 through -14; GM-CSF; adenosine deaminase; cellular growth factors, such as lymphokines; soluble CD4; Factor VIII; Factor IX; T-cell receptors; LDL receptor; ApoE; ApoC; alpha-1 antitrypsin; ornithine transcarbamylase (OTC); cystic fibrosis transmembrane receptor (CFTR); insulin; Fc receptors for antigen binding domains of antibodies, such as immunoglobulins; anti-HIV decoy tar elements;
15 and antisense sequences which inhibit viral replication, such as antisense sequences which inhibit replication of hepatitis B or hepatitis non-A, non-B virus. The nucleic acid is chosen considering several factors, including the cell to be transfected. Where the target cell is a blood cell, for example, particularly useful nucleic acids to use are those which allow the blood cells to exert a therapeutic effect, such as a gene encoding
20 a clotting factor for use in treatment of hemophilia. Another target cell is the lung airway cell, which can be used to administer nucleic acids, such as those coding for the cystic fibrosis transmembrane receptor, which could provide a gene therapeutic treatment for cystic fibrosis. Other target cells include muscle cells where useful nucleic acids, such as those encoding cytokines and growth factors, can be transduced
25 and the protein the nucleic acid encodes can be expressed and secreted to exert its effects on other cells, tissues and organs, such as the liver. Furthermore, the nucleic acid can encode more than one gene product, limited only, if the nucleic acid is to be packaged in a capsid, by the size of nucleic acid that can be packaged.

09717789.112100

Furthermore, suitable nucleic acids can include those that, when transferred into a primary cell, such as a blood cell, cause the transferred cell to target a site in the body where that cell's presence would be beneficial. For example, blood cells such as TIL cells can be modified, such as by transfer into the cell of a Fab portion of a monoclonal antibody, to recognize a selected antigen. Another example would be to introduce a nucleic acid that would target a therapeutic blood cell to tumor cells. Nucleic acids useful in treating cancer cells include those encoding chemotactic factors which cause an inflammatory response at a specific site, thereby having a therapeutic effect.

Cells, particularly blood cells, muscle cells, airway epithelial cells, brain cells and endothelial cells having such nucleic acids transferred into them can be useful in a variety of diseases, syndromes and conditions. For example, suitable nucleic acids include nucleic acids encoding soluble CD4, used in the treatment of AIDS and α -antitrypsin, used in the treatment of emphysema caused by α -antitrypsin deficiency.

Other diseases, syndromes and conditions in which such cells can be useful include, for example, adenosine deaminase deficiency, sickle cell deficiency, brain disorders such as Alzheimer's disease, thalassemia, hemophilia, diabetes, phenylketonuria, growth disorders and heart diseases, such as those caused by alterations in cholesterol metabolism, and defects of the immune system.

As another example, hepatocytes can be transfected with the present vectors having useful nucleic acids to treat liver disease. For example, a nucleic acid encoding OTC can be used to transfect hepatocytes (*ex vivo* and returned to the liver or *in vivo*) to treat congenital hyperammonemia, caused by an inherited deficiency in OTC. Another example is to use a nucleic acid encoding LDL to target hepatocytes *ex vivo* or *in vivo* to treat inherited LDL receptor deficiency. Such transfected hepatocytes can also be used to treat acquired infectious diseases, such as diseases resulting from a viral infection. For example, transduced hepatocyte precursors can be used to treat viral hepatitis, such as hepatitis B and non-A, non-B hepatitis, for example by transducing the hepatocyte precursor with a nucleic acid encoding an antisense RNA that inhibits

09717789 112100

viral replication. Another example includes transferring a vector of the present invention having a nucleic acid encoding a protein, such as α -interferon, which can confer resistance to the hepatitis virus.

5 For a procedure using transfected hepatocytes or hepatocyte precursors, hepatocyte precursors having a vector of the present invention transferred in can be grown in tissue culture, removed from the tissue culture vessel, and introduced to the body, such as by a surgical method. In this example, the tissue would be placed directly into the liver, or into the body cavity in proximity to the liver, as in a transplant
10 or graft. Alternatively, the cells can simply be directly injected into the liver, into the portal circulatory system, or into the spleen, from which the cells can be transported to the liver via the circulatory system. Furthermore, the cells can be attached to a support, such as microcarrier beads, which can then be introduced, such as by injection, into the peritoneal cavity. Once the cells are in the liver, by whatever means, the cells can then
15 express the nucleic acid and/or differentiate into mature hepatocytes which can express the nucleic acid.

The AAV5-derived vector can include any normally occurring AAV5 sequences in addition to an ITR and promoter. Examples of vector constructs are provided below.

20

The present vector or AAV5 particle or recombinant AAV5 virion can utilize any unique fragment of these present AAV5 nucleic acids, including the AAV5 nucleic acids set forth in SEQ ID NOS: 1 and 7-11, 13, 15, 16, 17, and 18. To be unique, the fragment must be of sufficient size to distinguish it from other known sequences, most
25 readily determined by comparing any nucleic acid fragment to the nucleotide sequences of nucleic acids in computer databases, such as GenBank. Such comparative searches are standard in the art. Typically, a unique fragment useful as a primer or probe will be at least about 8 or 10, preferable at least 20 or 25 nucleotides in length, depending upon the specific nucleotide content of the sequence. Additionally, fragments can be, for
30 example, at least about 30, 40, 50, 75, 100, 200 or 500 nucleotides in length and can

09747789.112100

encode polypeptides or be probes. The nucleic acid can be single or double stranded, depending upon the purpose for which it is intended. Where desired, the nucleic acid can be RNA.

5 The present invention further provides an AAV5 capsid protein to contain the vector. In particular, the present invention provides not only a polypeptide comprising all three AAV5 coat proteins, *i.e.*, VP1, VP2 and VP3, but also a polypeptide comprising each AAV5 coat protein individually, SEQ ID NOS: 4, 5, and 6, respectively. Thus an AAV5 particle comprising an AAV5 capsid protein comprises at
10 least one AAV5 coat protein VP1, VP2 or VP3. An AAV5 particle comprising an AAV5 capsid protein can be utilized to deliver a nucleic acid vector to a cell, tissue or subject. For example, the herein described AAV5 vectors can be encapsidated in an AAV5 capsid-derived particle and utilized in a gene delivery method. Furthermore, other viral nucleic acids can be encapsidated in the AAV5 particle and utilized in such
15 delivery methods. For example, an AAV1, 2,3,4,or 6 vector (e.g. AAV1,2,3,4,or 6 ITR and nucleic acid of interest)can be encapsidated in an AAV5 particle and administered. Furthermore, an AAV5 chimeric capsid incorporating both AAV2 capsid and AAV5 capsid sequences can be generated, by standard cloning methods, selecting regions from the known sequences of each protein as desired. For example, particularly
20 antigenic regions of the AAV2 capsid protein can be replaced with the corresponding region of the AAV5 capsid protein. In addition to chimeric capsids incorporating AAV2 capsid sequences, chimeric capsids incorporating AAV1, 3, 4, or 6 and AAV5 capsid sequences can be generated, by standard cloning methods, selecting regions from the known sequences of each protein as desired.

25

 The capsids can also be modified to alter their specific tropism by genetically altering the capsid to encode a specific ligand to a cell surface receptor. Alternatively, the capsid can be chemically modified by conjugating a ligand to a cell surface receptor. By genetically or chemically altering the capsids, the tropism can be
30 modified to direct AAV5 to a particular cell or population of cells. The capsids can

09717789-112100

also be altered immunologically by conjugating the capsid to an antibody that recognizes a specific protein on the target cell or population of cells.

The capsids can also be assembled into empty particles by expression in
5 mammalian, bacterial, fungal or insect cells. For example, AAV2 particles are known to be made from VP3 and VP2 capsid proteins in baculovirus. The same basic protocol can produce an empty AAV5 particle comprising an AAV5 capsid protein.

The herein described recombinant AAV5 nucleic acid derived vector can be
10 encapsidated in an AAV particle. In particular, it can be encapsidated in an AAV1 particle, an AAV2 particle, an AAV3 particle, an AAV4 particle, an AAV5 particle or an AAV6 particle, a portion of any of these capsids, or a chimeric capsid particle as described above, by standard methods using the appropriate capsid proteins in the encapsidation process, as long as the nucleic acid vector fits within the size limitation
15 of the particle utilized. The encapsidation process itself is standard in the art. The AAV5 replication machinery, i.e. the rep initiator proteins and other functions required for replication, can be utilized to produce the AAV5 genome that can be packaged in an AAV1, 2, 3, 4, 5 or 6 capsid.

20 The recombinant AAV5 virion containing a vector can also be produced by recombinant methods utilizing multiple plasmids. In one example, the AAV5 rep nucleic acid would be cloned into one plasmid, the AAV5 ITR nucleic acid would be cloned into another plasmid and the AAV1, 2, 3, 4, 5 or 6 capsid nucleic acid would be cloned on another plasmid. These plasmids would then be introduced into cells. The
25 cells that were efficiently transduced by all three plasmids, would exhibit specific integration as well as the ability to produce AAV5 recombinant virus. Additionally, two plasmids could be used where the AAV5 rep nucleic acid would be cloned into one plasmid and the AAV5 ITR and AAV5 capsid would be cloned into another plasmid. These plasmids would then be introduced into cells. The cells that were efficiently

09717789 112100

transduced by both plasmids, would exhibit specific integration as well as the ability to produce AAV5 recombinant virus.

An AAV5 capsid polypeptide encoding the entire VP1, VP2, and VP3 polypeptide can overall has greater than 56% homology to the polypeptide having the amino acid sequence encoded by nucleotides in SEQ ID NOS:7,8 and 9, as shown in figures 4 and 5. The capsid protein can have about 70% homology, about 75% homology, 80% homology, 85% homology, 90% homology, 95% homology, 98% homology, 99% homology, or even 100% homology to the protein having the amino acid sequence encoded by the nucleotides set forth in SEQ ID NOS:7, 8 or 9. The percent homology used to identify proteins herein, can be based on a nucleotide-by-nucleotide comparison or more preferable is based on a computerized algorithm as described herein. Variations in the amino acid sequence of the AAV5 capsid protein are contemplated herein, as long as the resulting particle comprising an AAV5 capsid protein remains antigenically or immunologically distinct from AAV1, AAV2, AAV3, AAV4 or AAV6 capsid, as can be routinely determined by standard methods. Specifically, for example, ELISA and Western blots can be used to determine whether a viral particle is antigenically or immunologically distinct from AAV2 or the other serotypes. Furthermore, the AAV5 particle preferably retains tissue tropism distinction from AAV2, such as that exemplified in the examples herein. An AAV5 chimeric particle comprising at least one AAV5 coat protein may have a different tissue tropism from that of an AAV5 particle consisting only of AAV5 coat proteins, but is still distinct from the tropism of an AAV2 particle.

The invention further provides a recombinant AAV5 virion, comprising an AAV5 particle containing, *i.e.*, encapsidating, a vector comprising a pair of AAV5 inverted terminal repeats. The recombinant vector can further comprise an AAV5 Rep-encoding nucleic acid. The vector encapsidated in the particle can further comprise an exogenous nucleic acid inserted between the inverted terminal repeats. AAV5 Rep confers targeted integration and efficient replication, thus production of recombinant

09717789 "112100

AAV5, comprising AAV5 Rep, yields more particles than production of recombinant AAV2. Since AAV5 is more efficient at replicating and packaging its genome, the exogenous nucleic acid inserted, or in the AAV5 capsids of the present invention, between the inverted terminal repeats can be packaged in the AAV1, 2, 3, 4, or 6 capsids to achieve the specific tissue tropism conferred by the capsid proteins.

The invention further contemplates chimeric recombinant ITRs that contains a rep binding site and a TRS site recognized by that Rep protein. By "Rep protein" is meant all four of the Rep proteins, Rep 40, Rep 78, Rep 52, Rep 68. Alternatively, "Rep protein" could be one or more of the Rep proteins described herein. One example of a chimeric ITR would consist of an AAV5 D region (SEQ ID NO: 23), an AAV5 TRS site (SEQ ID NO: 21), an AAV2 hairpin and an AAV2 binding site. Another example would be an AAV5 D region, an AAV5 TRS site, an AAV3 hairpin and an AAV3 binding site. In these chimeric ITRs, the D region can be from AAV1, 2, 3, 4, 5 or 6. The hairpin can be derived from AAV 1,2 3, 4, 5, 6. The binding site can be derived from any of AAV1, 2, 3, 4, 5 or 6. Preferably, the D region and the TRS are from the same serotype.

The chimeric ITRs can be combined with AAV5 Rep protein and any of the AAV serotype capsids to obtain recombinant virion. For example, recombinant virion can be produced by an AAV5 D region, an AAV5 TRS site, an AAV2 hairpin, an AAV2 binding site, AAV5 Rep protein and AAV1 capsid. This recombinant virion would possess the cellular tropism conferred by the AAV1 capsid protein and would possess the efficient replication conferred by the AAV5 Rep.

Other examples of the ITR, Rep protein and Capsids that will produce recombinant virus are provided in the list below:

5ITR + 5Rep + 5Cap=virus

5ITR + 5Rep + 1Cap=virus

- 5 ITR + 5Rep + 2Cap=virus
5 ITR + 5Rep + 3Cap=virus
5 ITR + 5Rep + 4Cap=virus
5 ITR + 5Rep + 6Cap=virus
5 1 ITR + 1Rep + 5Cap=virus
2 ITR + 2Rep + 5Cap=virus
3 ITR + 3Rep + 5Cap=virus
4 ITR + 4Rep + 5Cap=virus
6 ITR + 6Rep + 5Cap=virus

10

In any of the constructs described herein, inclusion of a promoter is preferred. As used in the constructs herein, unless otherwise specified, Cap (capsid) refers to any of AAV5 VP1, AAV5 VP2, AAV5 VP3, combinations thereof, functional fragments of any of VP1, VP2 or VP3, or chimeric capsids as described herein. The ITRs of the
15 constructs described herein, can be chimeric recombinant ITRs as described elsewhere in the application.

Conjugates of recombinant or wild-type AAV5 virions and nucleic acids or proteins can be used to deliver those molecules to a cell. For example, the purified
20 AAV5 can be used as a vehicle for delivering DNA bound to the exterior of the virus. Examples of this are to conjugate the DNA to the virion by a bridge using poly-L-lysine or other charged molecule. Also contemplated are virosomes that contain AAV5 structural proteins (AAV5 capsid proteins), lipids such as DOTAP, and nucleic acids that are complexed via charge interaction to introduce DNA into cells.

25

Also provided by this invention are conjugates that utilize the AAV5 capsid or a unique region of the AAV5 capsid protein (e.g. VP1, VP2 or VP3 or combinations thereof) to introduce DNA into cells. For example, the type 5 VP3 protein or fragment thereof, can be conjugated to a DNA on a plasmid that is conjugated to a lipid. Cells
30 can be infected using the targeting ability of the VP3 capsid protein to achieve the

09717789 "112100

desired tissue tropism, specific to AAV5. Type 5 VP1 and VP2 proteins can also be utilized to introduce DNA or other molecules into cells. By further incorporating the Rep protein and the AAV TRS into the DNA-containing conjugate, cells can be transduced and targeted integration can be achieved. For example, if AAV5 specific
5 targeted integration is desired, a conjugate composed of the AAV5 VP3 capsid, AAV5 rep or a fragment of AAV5 rep, AAV5 TRS, the rep binding site, the heterologous DNA of interest, and a lipid, can be utilized to achieve AAV5 specific tropism and AAV5 specific targeted integration in the genome.

10 Further provided by this invention are chimeric viruses where AAV5 can be combined with herpes virus, baculovirus or other viruses to achieve a desired tropism associated with another virus. For example, the AAV5 ITRs could be inserted in the herpes virus and cells could be infected. Post-infection, the ITRs of AAV5 could be acted on by AAV5 rep provided in the system or in a separate vehicle to rescue AAV5
15 from the genome. Therefore, the cellular tropism of the herpes simplex virus can be combined with AAV5 rep mediated targeted integration. Other viruses that could be utilized to construct chimeric viruses include, lentivirus, retrovirus, pseudotyped retroviral vectors, and adenoviral vectors.

20 The present invention further provides isolated nucleic acids of AAV5. For example, provided is an isolated nucleic acid comprising the nucleotide sequence set forth in SEQ ID NO:1 (AAV5 genome). This nucleic acid, or portions thereof, can be inserted into vectors, such as plasmids, yeast artificial chromosomes, or other viral vector (particle), if desired, by standard cloning methods. The present invention also
25 provides an isolated nucleic acid consisting essentially of the nucleotide sequence set forth in SEQ ID NO:1. The nucleotides of SEQ ID NO:1 can have minor modifications and still be contemplated by the present invention. For example, modifications that do not alter the amino acid encoded by any given codon (such as by modification of the third, "wobble," position in a codon) can readily be made, and such alterations are
30 known in the art. Furthermore, modifications that cause a resulting neutral (conserved)

09717789 "112100

amino acid substitution of a similar amino acid can be made in a coding region of the genome. Additionally, modifications as described herein for the AAV5 components, such as the ITRs, the p5 promoter, etc. are contemplated in this invention.

Furthermore, modifications to regions of SEQ ID NO:1 other than in the ITR, TRS Rep
5 binding site and hairpin are likely to be tolerated without serious impact on the function of the nucleic acid as a recombinant vector.

As used herein, the term "isolated" refers to a nucleic acid separated or significantly free from at least some of the other components of the naturally occurring
10 organism, for example, the cell structural components or viral components commonly found associated with nucleic acids in the environment of the virus and/or other nucleic acids. The isolation of the native nucleic acids can be accomplished, for example, by techniques such as cell lysis followed by phenol plus chloroform extraction, followed by ethanol precipitation of the nucleic acids. The nucleic acids of this invention can be
15 isolated from cells according to any of many methods well known in the art.

As used herein, the term "nucleic acid" refers to single-or multiple stranded molecules which may be DNA or RNA, or any combination thereof, including modifications to those nucleic acids. The nucleic acid may represent a coding strand or
20 its complement, or any combination thereof. Nucleic acids may be identical in sequence to the sequences which are naturally occurring for any of the novel genes discussed herein or may include alternative codons which encode the same amino acid as those provided herein, including that which is found in the naturally occurring sequence. These nucleic acids can also be modified from their typical structure. Such
25 modifications include, but are not limited to, methylated nucleic acids, the substitution of a non-bridging oxygen on the phosphate residue with either a sulfur (yielding phosphorothioate deoxynucleotides), selenium (yielding phosphorselenoate deoxynucleotides), or methyl groups (yielding methylphosphonate deoxynucleotides).

09717789 112100

09717789 112100

The present invention additionally provides an isolated nucleic acid that selectively hybridizes with any nucleic acid disclosed herein, including the entire AAV5 genome and any unique fragment thereof, including the Rep and capsid encoding sequences (e.g. SEQ ID NOS: 1, 7, 8, 9, 10, 11, 13, 15, 16, 17, 18, 19, 20, 21, 22 and 23). Specifically, the nucleic acid can selectively or specifically hybridize to an isolated nucleic acid consisting of the nucleotide sequence set forth in SEQ ID NO:1 (AAV5 genome). The present invention further provides an isolated nucleic acid that selectively or specifically hybridizes with an isolated nucleic acid comprising the nucleotide sequence set forth in SEQ ID NO:1 (AAV5 genome). By "selectively hybridizes" as used herein is meant a nucleic acid that hybridizes to one of the disclosed nucleic acids under sufficient stringency conditions without significant hybridization to a nucleic acid encoding an unrelated protein, and particularly, without detectably hybridizing to nucleic acids of AAV2. Thus, a nucleic acid that selectively hybridizes with a nucleic acid of the present invention will not selectively hybridize under stringent conditions with a nucleic acid encoding a different protein or the corresponding protein from a different serotype of the virus, and vice versa. A "specifically hybridizing" nucleic acid is one that hybridizes under stringent conditions to only a nucleic acid found in AAV5. Therefore, nucleic acids for use, for example, as primers and probes to detect or amplify the target nucleic acids are contemplated herein. Nucleic acid fragments that selectively hybridize to any given nucleic acid can be used, e.g., as primers and or probes for further hybridization or for amplification methods (e.g., polymerase chain reaction (PCR), ligase chain reaction (LCR)). Additionally, for example, a primer or probe can be designed that selectively hybridizes with both AAV5 and a gene of interest carried within the AAV5 vector (i.e., a chimeric nucleic acid).

Stringency of hybridization is controlled by both temperature and salt concentration of either or both of the hybridization and washing steps. Typically, the stringency of hybridization to achieve selective hybridization involves hybridization in high ionic strength solution (6X SSC or 6X SSPE) at a temperature that is about 12-

25°C below the T_m (the melting temperature at which half of the molecules dissociate from their hybridization partners) followed by washing at a combination of temperature and salt concentration chosen so that the washing temperature is about 5°C to 20°C below the T_m . The temperature and salt conditions are readily determined empirically
5 in preliminary experiments in which samples of reference DNA immobilized on filters are hybridized to a labeled nucleic acid of interest and then washed under conditions of different stringencies. Hybridization temperatures are typically higher for DNA-RNA and RNA-RNA hybridizations. The washing temperatures can be used as described above to achieve selective stringency, as is known in the art. (Sambrook et al.,
10 *Molecular Cloning: A Laboratory Manual*, 2nd Ed., Cold Spring Harbor Laboratory, Cold Spring Harbor, New York, 1989; Kunkel et al. *Methods Enzymol.* 1987:154:367, 1987). A preferable stringent hybridization condition for a DNA:DNA hybridization can be at about 68°C (in aqueous solution) in 6X SSC or 6X SSPE followed by washing at 68°C. Stringency of hybridization and washing, if desired, can be reduced
15 accordingly as the degree of complementarity desired is decreased, and further, depending upon the G-C or A-T richness of any area wherein variability is searched for. Likewise, stringency of hybridization and washing, if desired, can be increased accordingly as homology desired is increased, and further, depending upon the G-C or A-T richness of any area wherein high homology is desired, all as known in the art.

20

A nucleic acid that selectively hybridizes to any portion of the AAV5 genome is contemplated herein. Therefore, a nucleic acid that selectively hybridizes to AAV5 can be of longer length than the AAV5 genome, it can be about the same length as the AAV5 genome or it can be shorter than the AAV5 genome. The length of the nucleic
25 acid is limited on the shorter end of the size range only by its specificity for hybridization to AAV5, *i.e.*, once it is too short, typically less than about 5 to 7 nucleotides in length, it will no longer bind specifically to AAV5, but rather will hybridize to numerous background nucleic acids. Additionally contemplated by this invention is a nucleic acid that has a portion that specifically hybridizes to AAV5 and a
30 portion that specifically hybridizes to a gene of interest inserted within AAV5.

09717789-112100

09717789 112100

The present invention further provides an isolated nucleic acid encoding an adeno-associated virus 5 Rep protein. The AAV5 Rep proteins are encoded by open reading frame (ORF) 1 of the AAV5 genome. Examples of the AAV5 Rep genes are shown in the nucleic acid set forth in SEQ ID NO:1, and include nucleic acids

5 consisting essentially of the nucleotide sequences set forth in SEQ ID NOS:10 (Rep52), 11 (Rep78), 13 (Rep40), and 15 (Rep68), and nucleic acids comprising the nucleotide sequences set forth in SEQ ID NOS:10 , 11, 13, and 15. However, the present invention contemplates that the Rep nucleic acid can include any one, two, three, or four of the four Rep proteins, in any order, in such a nucleic acid. Furthermore, minor

10 modifications are contemplated in the nucleic acid, such as silent mutations in the coding sequences, mutations that make neutral or conservative changes in the encoded amino acid sequence, and mutations in regulatory regions that do not disrupt the expression of the gene. Examples of other minor modifications are known in the art. Further modifications can be made in the nucleic acid, such as to disrupt or alter

15 expression of one or more of the Rep proteins in order to, for example, determine the effect of such a disruption; such as to mutate one or more of the Rep proteins to determine the resulting effect, etc. However, in general, a modified nucleic acid encoding a Rep protein will have at least about 85%, about 90%, about 93%, about 95%, about 98% or 100% homology to the Rep nucleic sequences described herein

20 e.g., SEQ ID NOS: 10, 11, 13 and 15, and the Rep polypeptide encoded therein will have overall about 93%, about 95%, about 98%, about 99% or 100% homology with the amino acid sequence described herein, e.g., SEQ ID NOS:2 , 3, 12 and 14. Percent homology is determined by the techniques described herein.

25 The present invention also provides an isolated nucleic acid that selectively or specifically hybridizes with a nucleic acid consisting essentially of the nucleotide sequence set forth in SEQ ID NOS:10, 11, 13 and 15, and an isolated nucleic acid that selectively hybridizes with a nucleic acid comprising the nucleotide sequence set forth in SEQ ID NOS:10, 11, 13 and 15. "Selectively hybridizing" and "stringency of

30 hybridization" is defined elsewhere herein.

As described above, the present invention provides the nucleic acid encoding a Rep 40 protein and, in particular an isolated nucleic acid comprising the nucleotide sequence set forth in SEQ ID NO: 13, an isolated nucleic acid consisting essentially of the nucleotide sequence set forth in SEQ ID NO: 13, and a nucleic acid encoding the adeno-associated virus 5 protein having the amino acid sequence set forth in SEQ ID NO: 12. The present invention also provides the nucleic acid encoding a Rep 52 protein, and in particular an isolated nucleic acid comprising the nucleotide sequence set forth in SEQ ID NO:10, an isolated nucleic acid consisting essentially of the nucleotide sequence set forth in SEQ ID NO:10, and a nucleic acid encoding the adeno-associated virus 5 Rep protein having the amino acid sequence set forth in SEQ ID NO:2. The present invention further provides the nucleic acid encoding a Rep 68 protein and, in particular an isolated nucleic acid comprising the nucleotide sequence set forth in SEQ ID NO: 15, an isolated nucleic acid consisting essentially of the nucleotide sequence set forth in SEQ ID NO: 15, and a nucleic acid encoding the adeno-associated virus 5 protein having the amino acid sequence set forth in SEQ ID NO: 14. And, further, the present invention provides the nucleic acid encoding a Rep 78 protein, and in particular an isolated nucleic acid comprising the nucleotide sequence set forth in SEQ ID NO:11, an isolated nucleic acid consisting essentially of the nucleotide sequence set forth in SEQ ID NO:11, and a nucleic acid encoding the adeno-associated virus 5 Rep protein having the amino acid sequence set forth in SEQ ID NO:3. As described elsewhere herein, these nucleic acids can have minor modifications, including silent nucleotide substitutions, mutations causing conservative amino acid substitutions in the encoded proteins, and mutations in control regions that do not or minimally affect the encoded amino acid sequence.

25

The present invention further provides a nucleic acid encoding the entire AAV5 Capsid polypeptide. Furthermore, the present invention provides a nucleic acid encoding each of the three AAV5 coat proteins, VP1, VP2, and VP3. Thus, the present invention provides a nucleic acid encoding AAV5 VP1, a nucleic acid encoding AAV5 VP2, and a nucleic acid encoding AAV5 VP3. Thus, the present invention provides a

09717789 112100
001211 6827260

09747789 "112100

nucleic acid encoding the amino acid sequence set forth in SEQ ID NO:4 (VP1); a nucleic acid encoding the amino acid sequence set forth in SEQ ID NO:5 (VP2), and a nucleic acid encoding the amino acid sequence set forth in SEQ ID NO:6 (VP3). The present invention also specifically provides a nucleic acid comprising SEQ ID NO:7 (VP1 gene); a nucleic acid comprising SEQ ID NO:8 (VP2 gene); and a nucleic acid comprising SEQ ID NO:9 (VP3 gene). The present invention also specifically provides a nucleic acid consisting essentially of SEQ ID NO:7 (VP1 gene), a nucleic acid consisting essentially of SEQ ID NO:8 (VP2 gene), and a nucleic acid consisting essentially of SEQ ID NO:9 (VP3 gene). Minor modifications in the nucleotide sequences encoding the capsid, or coat, proteins are contemplated, as described above for other AAV5 nucleic acids. However, in general, a modified nucleic acid encoding a capsid protein will have at least about 85%, about 90%, about 93%, about 95%, about 98% or 100% homology to the capsid nucleic sequences described herein e.g., SEQ ID NOS: 7, 8, and 9, and the capsid polypeptide encoded therein will have overall about 93%, about 95%, about 98%, about 99% or 100% homology with the amino acid sequence described herein, e.g., SEQ ID NOS:4, 5, and 6. Nucleic acids that selectively hybridize with the nucleic acids of SEQ ID NOS:7,8 and 9 under the conditions described above are also provided.

20 The present invention also provides a cell containing one or more of the herein described nucleic acids, such as the AAV5 genome, AAV5 ORF1 and ORF2, each AAV5 Rep protein gene, or each AAV5 capsid protein gene. Such a cell can be any desired cell and can be selected based upon the use intended. For example, cells can include bacterial cells, yeast cells, insect cells, human HeLa cells and simian Cos cells as well as other human and mammalian cells and cell lines. Primary cultures as well as established cultures and cell lines can be used. Nucleic acids of the present invention can be delivered into cells by any selected means, in particular depending upon the target cells. Many delivery means are well-known in the art. For example, electroporation, calcium phosphate precipitation, microinjection, cationic or anionic liposomes, and liposomes in combination with a nuclear localization signal peptide for

30

delivery to the nucleus can be utilized, as is known in the art. Additionally, if the nucleic acids are in a viral particle, the cells can simply be transduced with the virion by standard means known in the art for AAV transduction. Small amounts of the recombinant AAV5 virus can be made to infect cells and produce more of itself.

5

The invention provides purified AAV5 polypeptides. The term "polypeptide" as used herein refers to a polymer of amino acids and includes full-length proteins and fragments thereof. Thus, "protein," polypeptide," and "peptide" are often used interchangeably herein. Substitutions can be selected by known parameters to be neutral (*see, e.g.*, Robinson WE Jr, and Mitchell WM., AIDS 4:S151-S162 (1990)). As will be appreciated by those skilled in the art, the invention also includes those polypeptides having slight variations in amino acid sequences or other properties. Such variations may arise naturally as allelic variations (*e.g.*, due to genetic polymorphism) or may be produced by human intervention (*e.g.*, by mutagenesis of cloned DNA sequences), such as induced point, deletion, insertion and substitution mutants. Minor changes in amino acid sequence are generally preferred, such as conservative amino acid replacements, small internal deletions or insertions, and additions or deletions at the ends of the molecules. Substitutions may be designed based on, for example, the model of Dayhoff, *et al.* (in *Atlas of Protein Sequence and Structure* 1978, Nat'l Biomed. Res. Found., Washington, D.C.). These modifications can result in changes in the amino acid sequence, provide silent mutations, modify a restriction site, or provide other specific mutations. The location of any modifications to the polypeptide will often determine its impact on function. Particularly, alterations in regions non-essential to protein function will be tolerated with fewer effects on function. Elsewhere in the application regions of the AAV5 proteins are described to provide guidance as to where substitutions, additions or deletions can be made to minimize the likelihood of disturbing the function of the variant.

A polypeptide of the present invention can be readily obtained by any of several means. For example, the polypeptide of interest can be synthesized chemically by

09717789-112100

standard methods. Additionally, the coding regions of the genes can be recombinantly expressed and the resulting polypeptide isolated by standard methods. Furthermore, an antibody specific for the resulting polypeptide can be raised by standard methods (see, e.g., Harlow and Lane, *Antibodies: A Laboratory Manual*, Cold Spring Harbor Laboratory, Cold Spring Harbor, New York, 1988), and the protein can be isolated from a cell expressing the nucleic acid encoding the polypeptide by selective hybridization with the antibody. This protein can be purified to the extent desired by standard methods of protein purification (see, e.g., Sambrook et al., *Molecular Cloning: A Laboratory Manual*, 2nd Ed., Cold Spring Harbor Laboratory, Cold Spring Harbor, New York, 1989).

Typically, to be unique, a polypeptide fragment of the present invention will be at least about 5 amino acids in length; however, unique fragments can be 6, 7, 8, 9, 10, 20, 30, 40, 50, 60, 70, 80, 90, 100 or more amino acids in length. A unique polypeptide will typically comprise such a unique fragment; however, a unique polypeptide can also be determined by its overall homology. A unique polypeptide can be 6, 7, 8, 9, 10, 20, 30, 40, 50, 60, 70, 80, 90, 100 or more amino acids in length. Uniqueness of a polypeptide fragment can readily be determined by standard methods such as searches of computer databases of known peptide or nucleic acid sequences or by hybridization studies to the nucleic acid encoding the protein or to the protein itself, as known in the art. The uniqueness of a polypeptide fragment can also be determined immunologically as well as functionally. Uniqueness can be simply determined in an amino acid-by-amino acid comparison of the polypeptides.

An antigenic or immunoreactive fragment of this invention is typically an amino acid sequence of at least about 5 consecutive amino acids, and it can be derived from the AAV5 polypeptide amino acid sequence. An antigenic AAV5 fragment is any fragment unique to the AAV5 protein, as described herein, against which an AAV5-specific antibody can be raised, by standard methods. Thus, the resulting antibody-antigen reaction should be specific for AAV5.

09717789-112100

09717789-112100

The present invention provides an isolated AAV5 Rep protein. An AAV5 Rep polypeptide is encoded by ORF1 of AAV5. The present invention also provides each individual AAV5 Rep protein. Thus the present invention provides AAV5 Rep 40 (e.g., SEQ ID NO: 12), or a unique fragment thereof. The present invention provides
5 AAV5 Rep 52 (e.g., SEQ ID NO: 2), or a unique fragment thereof. The present invention provides AAV5 Rep 68 (e.g., SEQ ID NO: 14), or a unique fragment thereof. The present invention provides an example of AAV5 Rep 78 (e.g., SEQ ID NO: 3), or a unique fragment thereof. By "unique fragment thereof" is meant any smaller polypeptide fragment encoded by an AAV5 rep gene that is of sufficient length to be
10 found only in the Rep polypeptide. Substitutions and modifications of the amino acid sequence can be made as described above and, further, can include protein processing modifications, such as glycosylation, to the polypeptide.

The present invention further provides an AAV5 Capsid polypeptide or a
15 unique fragment thereof. AAV5 capsid polypeptide is encoded by ORF 2 of AAV5. The present invention further provides the individual AAV5 capsid proteins, VP1, VP2 and VP3 or unique fragments thereof. Thus, the present invention provides an isolated polypeptide having the amino acid sequence set forth in SEQ ID NO:4 (VP1). The present invention additionally provides an isolated polypeptide having the amino acid
20 sequence set forth in SEQ ID NO:5 (VP2). The present invention also provides an isolated polypeptide having the amino acid sequence set forth in SEQ ID NO:6 (VP3). By "unique fragment thereof" is meant any smaller polypeptide fragment encoded by any AAV5 capsid gene that is of sufficient length to be found only in the AAV5 capsid protein. Substitutions and modifications of the amino acid sequence can be made as
25 described above and, further, can include protein processing modifications, such as glycosylation, to the polypeptide. However, an AAV5 Capsid polypeptide including all three coat proteins will have greater than about 56% overall homology to the polypeptide encoded by the nucleotides set forth in SEQ ID NOS:4,5 or 6. The protein can have about 65%, about 70%, about 75%, about 80%, about 85%, about 90%, 93%,
30 95%, 97% or even 100% homology to the amino acid sequence encoded by the

nucleotides set forth in SEQ ID NOS:4,5 or 6. An AAV5 VP1 polypeptide can have at least about 58%, about 60%, about 70%, about 80%, about 90%, 93%, 95%, 97% or about 100% homology to the amino acid sequence set forth in SEQ ID NO:4. An AAV5 VP2 polypeptide can have at least about 58%, about 60%, about 70%, about 80%, about 90%, 93%, 95%, 97% or about 100% homology to the amino acid sequence set forth in SEQ ID NO:5. An AAV5 VP3 polypeptide can have at least about 60%, about 70%, about 80%, about 90%, 93%, 95%, 97% or about 100% homology to the amino acid sequence set forth in SEQ ID NO:6.

10 The present invention further provides an isolated antibody that specifically binds an AAV5 Rep protein or a unique epitope thereof. Also provided are isolated antibodies that specifically bind the AAV5 Rep 52 protein, the AAV5 Rep 40 protein, the AAV5 Rep 68 protein and the AAV5 Rep 78 protein having the amino acid sequences set forth in SEQ ID NO:2, SEQ ID NO: 12, SEQ ID NO: 14 and SEQ ID NO: 3, respectively or that specifically binds a unique fragment thereof. Clearly, any given antibody can recognize and bind one of a number of possible epitopes present in the polypeptide; thus only a unique portion of a polypeptide (having the epitope) may need to be present in an assay to determine if the antibody specifically binds the polypeptide.

20 The present invention additionally provides an isolated antibody that specifically binds any of the adeno-associated virus 5 Capsid proteins (VP1, VP2 or VP3), a unique epitope thereof, or the polypeptide comprising all three AAV5 coat proteins. Also provided is an isolated antibody that specifically binds the AAV5 capsid protein having the amino acid sequence set forth in SEQ ID NO:4 (VP1), or that specifically binds a unique fragment thereof. The present invention further provides an isolated antibody that specifically binds the AAV5 Capsid protein having the amino acid sequence set forth in SEQ ID NO:5 (VP2), or that specifically binds a unique fragment thereof. The invention additionally provides an isolated antibody that specifically binds the AAV5 Capsid protein having the amino acid sequence set forth in

09717789.112100

SEQ ID NO:6 (VP3), or that specifically binds a unique fragment thereof. Again, any given antibody can recognize and bind one of a number of possible epitopes present in the polypeptide; thus only a unique portion of a polypeptide (having the epitope) may need to be present in an assay to determine if the antibody specifically binds the
5 polypeptide.

The antibody can be a component of a composition that comprises an antibody that specifically binds the AAV5 protein. The composition can further comprise, *e.g.*, serum, serum-free medium, or a pharmaceutically acceptable carrier such as
10 physiological saline, etc..

By "an antibody that specifically binds" an AAV5 polypeptide or protein is meant an antibody that selectively binds to an epitope on any portion of the AAV5 peptide such that the antibody binds specifically to the corresponding AAV5
15 polypeptide without significant background. Specific binding by an antibody further means that the antibody can be used to selectively remove the target polypeptide from a sample comprising the polypeptide or and can readily be determined by radioimmunoassay (RIA), bioassay, or enzyme-linked immunosorbant (ELISA) technology. An ELISA method effective for the detection of the specific antibody-
20 antigen binding can, for example, be as follows: (1) bind the antibody to a substrate; (2) contact the bound antibody with a sample containing the antigen; (3) contact the above with a secondary antibody bound to a detectable moiety (*e.g.*, horseradish peroxidase enzyme or alkaline phosphatase enzyme); (4) contact the above with the substrate for the enzyme; (5) contact the above with a color reagent; (6) observe the
25 color change.

An antibody can include antibody fragments such as Fab fragments which retain the binding activity. Antibodies can be made as described in, *e.g.*, Harlow and Lane, *Antibodies: A Laboratory Manual*, Cold Spring Harbor Laboratory, Cold Spring
30 Harbor, New York (1988). Briefly, purified antigen can be injected into an animal in

09717789 112100

an amount and in intervals sufficient to elicit an immune response. Antibodies can either be purified directly, or spleen cells can be obtained from the animal. The cells are then fused with an immortal cell line and screened for antibody secretion. Individual hybridomas are then propagated as individual clones serving as a source for
5 a particular monoclonal antibody.

The present invention additionally provides a method of screening a cell for infectivity by AAV5 comprising contacting the cell with AAV5 and detecting the presence of AAV5 in the cells. AAV5 particles can be detected using any standard physical or biochemical methods. For example, physical methods that can be used for
10 this detection include DNA based methods such as 1) polymérase chain reaction (PCR) for viral DNA or RNA or 2) direct hybridization with labeled probes, and immunological methods such as by 3) antibody directed against the viral structural or non- structural proteins. Catalytic methods of viral detection include, but are not limited to, detection of site and strand specific DNA nicking activity of Rep proteins or
15 replication of an AAV origin- containing substrate. Reporter genes can also be utilized to detect cells that transduct AAV-5. For example, β -gal, green flourescent protein or luciferase can be inserted into a recombinant AAV-5. The cell can then be contacted with the recombinant AAV-5, either *in vitro* or *in vivo* and a colorimetric assay could detect a color change in the cells that would indicate transduction of AAV-5 in the cell.
20 Additional detection methods are outlined in Fields, *Virology*, Raven Press, New York, New York. 1996.

For screening a cell for infectivity by AAV5, wherein the presence of AAV5 in the cells is determined by nucleic acid hybridization methods, a nucleic acid probe for
25 such detection can comprise, for example, a unique fragment of any of the AAV5 nucleic acids provided herein. The uniqueness of any nucleic acid probe can readily be determined as described herein. Additionally, the presence of AAV5 in cells can be determined by flourescence, antibodies to gene products, focus forming assays, plaque lifts, Western blots and chromogenic assays. The nucleic acid can be, for example, the

09717789.112100

nucleic acid whose nucleotide sequence is set forth in SEQ ID NO: 1,7, 8, 9, 10, 11, 13, 15, 16, 17, 18, 19, 20, 21, 22, 23 or a unique fragment thereof.

09717789-112100

The present invention includes a method of determining the suitability of an AAV5 vector for administration to a subject comprising administering to an antibody-containing sample from the subject an antigenic fragment of an isolated AAV5 Rep or Capsid protein, and detecting neutralizing antibody-antigen reaction in the sample, the presence of a neutralizing reaction indicating the AAV5 vector may be unsuitable for use in the subject. The present method of determining the suitability of an AAV5 vector for administration to a subject can comprise contacting an antibody-containing sample from the subject with a unique antigenic or immunogenic fragment of an AAV5 Rep protein (e.g. Rep 40, Rep 52, Rep 68, Rep 78) and detecting an antibody-antigen reaction in the sample, the presence of a reaction indicating the AAV5 vector to be unsuitable for use in the subject. The AAV5 Rep proteins are provided herein, and their antigenic fragments are routinely determined. The AAV5 capsid protein can be used to select an antigenic or immunogenic fragment, for example from the amino acid sequence set forth in SEQ ID NO:4 (VP1), the amino acid sequence set forth in SEQ ID NO: 5 (VP2) or the amino acid sequence set forth in SEQ ID NO:6 (VP3). Alternatively, or additionally, an antigenic or immunogenic fragment of an isolated AAV5 Rep protein can be utilized in this determination method. The AAV5 Rep protein from which an antigenic fragment is selected can have the amino acid sequence encoded by the nucleic acid set forth in SEQ ID NO:1, the amino acid sequence set forth in SEQ ID NO:2, or the amino acid sequence set forth in SEQ ID NO:3, the amino acid sequence set forth in SEQ ID NO: 12, or the amino acid sequence set forth in SEQ ID NO:14.

The AAV5 polypeptide fragments can be analyzed to determine their antigenicity, immunogenicity and/or specificity. Briefly, various concentrations of a putative immunogenically specific fragment are prepared and administered to a subject and the immunological response (e.g., the production of antibodies or cell mediated

immunity) of an animal to each concentration is determined. The amounts of antigen administered depend on the subject, e.g. a human, rabbit or a guinea pig, the condition of the subject, the size of the subject, etc. Thereafter an animal so inoculated with the antigen can be exposed to the AAV5 viral particle or AAV5 protein to test the

5 immunoreactivity or the antigenicity of the specific immunogenic fragment. The specificity of a putative antigenic or immunogenic fragment can be ascertained by testing sera, other fluids or lymphocytes from the inoculated animal for cross reactivity with other closely related viruses, such as AAV1, AAV2, AAV3, AAV4 and AAV5.

10 The hemagglutination assay can also be used to rapidly identify and detect AAV5 viral particles. Detection of hemagglutination activity correlates with infectivity and can be used to titer the virus. This assay could also be used to identify antibodies in a patients serum which might interact with the virus. Hemagglutination has been shown to correlate with infectivity and therefore hemagglutination may be a useful
15 assay for identify cellular receptors for AAV5.

By the "suitability of an AAV5 vector for administration to a subject" is meant a determination of whether the AAV5 vector will elicit a neutralizing immune response upon administration to a particular subject. A vector that does not elicit a significant
20 immune response is a potentially suitable vector, whereas a vector that elicits a significant, neutralizing immune response (e.g. at least 90%) is thus likely to be unsuitable for use in that subject. Significance of any detectable immune response is a standard parameter understood by the skilled artisan in the field. For example, one can incubate the subject's serum with the virus, then determine whether that virus retains its
25 ability to transduce cells in culture. If such virus cannot transduce cells in culture, the vector likely has elicited a significant immune response.

Alternatively, or additionally, one skilled in the art could determine whether or not AAV5 administration would be suitable for a particular cell type of a subject. For
30 example, the artisan could culture muscle cells *in vitro* and transduce the cells with

09717789 "112100

AAV5 in the presence or absence of the subject's serum. If there is a reduction in transduction efficiency, this could indicate the presence of a neutralizing antibody or other factors that may inhibit transduction. Normally, greater than 90% inhibition would have to be observed in order to rule out the use of AAV-5 as a vector. However,
5 this limitation could be overcome by treating the subject with an immunosuppressant that could block the factors inhibiting transduction.

As will be recognized by those skilled in the art, numerous types of immunoassays are available for use in the present invention to detect binding between
10 an antibody and an AAV5 polypeptide of this invention. For instance, direct and indirect binding assays, competitive assays, sandwich assays, and the like, as are generally described in, e.g., U.S. Pat. Nos. 4,642,285; 4,376,110; 4,016,043; 3,879,262; 3,852,157; 3,850,752; 3,839,153; 3,791,932; and Harlow and Lane, *Antibodies, A Laboratory Manual*, Cold Spring Harbor Publications, N.Y. (1988). For example,
15 enzyme immunoassays such as immunofluorescence assays (IFA), enzyme linked immunosorbent assays (ELISA) and immunoblotting can be readily adapted to accomplish the detection of the antibody. An ELISA method effective for the detection of the antibody bound to the antigen can, for example, be as follows: (1) bind the antigen to a substrate; (2) contact the bound antigen with a fluid or tissue sample
20 containing the antibody; (3) contact the above with a secondary antibody specific for the antigen and bound to a detectable moiety (e.g., horseradish peroxidase enzyme or alkaline phosphatase enzyme); (4) contact the above with the substrate for the enzyme; (5) contact the above with a color reagent; (6) observe color change.

25 The antibody-containing sample of this method can comprise any biological sample which would contain the antibody or a cell containing the antibody, such as blood, plasma, serum, bone marrow, saliva and urine.

The present invention also provides a method of producing the AAV5 virus by
30 transducing a cell with the nucleic acid encoding the virus.

09717789 "112100

The present method further provides a method of delivering an exogenous (heterologous) nucleic acid to a cell comprising administering to the cell an AAV5 particle containing a vector comprising the nucleic acid inserted between a pair of AAV inverted terminal repeats, thereby delivering the nucleic acid to the cell.

5

The AAV ITRs in the vector for the herein described delivery methods can be AAV5 ITRs (SEQ ID NOS: 19 and 20). Furthermore, the AAV ITRs in the vector for the herein described nucleic acid delivery methods can also comprise AAV1, AAV2, AAV3, AAV4, or AAV6 inverted terminal repeats.

10

The present invention also includes a method of delivering a heterologous nucleic acid to a subject comprising administering to a cell from the subject an AAV5 particle containing a vector comprising the nucleic acid inserted between a pair of AAV inverted terminal repeats, and returning the cell to the subject, thereby delivering the nucleic acid to the subject. The AAV ITRs can be any AAV ITRs, including AAV5 ITRs and AAV2 ITRs. For example, in an *ex vivo* administration, cells are isolated from a subject by standard means according to the cell type and placed in appropriate culture medium, again according to cell type (*see, e.g.*, ATCC catalog). Viral particles are then contacted with the cells as described above, and the virus is allowed to transduce the cells. Cells can then be transplanted back into the subject's body, again by means standard for the cell type and tissue (*e. g.*, in general, U.S. Patent No. 5,399,346; for neural cells, Dunnett, S.B. and Björklund, A., eds., *Transplantation: Neural Transplantation-A Practical Approach*, Oxford University Press, Oxford (1992)). If desired, prior to transplantation, the cells can be studied for degree of transduction by the virus, by known detection means and as described herein. Cells for *ex vivo* transduction followed by transplantation into a subject can be selected from those listed above, or can be any other selected cell. Preferably, a selected cell type is examined for its capability to be transfected by AAV5. Preferably, the selected cell will be a cell readily transduced with AAV5 particles; however, depending upon the application, even cells with relatively low transduction efficiencies can be useful,

30

09747789-112100

particularly if the cell is from a tissue or organ in which even production of a small amount of the protein or antisense RNA encoded by the vector will be beneficial to the subject.

5 The present invention further provides a method of delivering a nucleic acid to a cell in a subject comprising administering to the subject an AAV5 particle containing a vector comprising the nucleic acid inserted between a pair of AAV inverted terminal repeats, thereby delivering the nucleic acid to a cell in the subject. Administration can be an *ex vivo* administration directly to a cell removed from a subject, such as any of
10 the cells listed above, followed by replacement of the cell back into the subject, or administration can be *in vivo* administration to a cell in the subject. For *ex vivo* administration, cells are isolated from a subject by standard means according to the cell type and placed in appropriate culture medium, again according to cell type (*see, e.g.,* ATCC catalog). Viral particles are then contacted with the cells as described above,
15 and the virus is allowed to transfect the cells. Cells can then be transplanted back into the subject's body, again by means standard for the cell type and tissue (*e. g.,* for neural cells, Dunnett, S.B. and Björklund, A., eds., *Transplantation: Neural Transplantation-A Practical Approach*, Oxford University Press, Oxford (1992)). If desired, prior to transplantation, the cells can be studied for degree of transfection by
20 the virus, by known detection means and as described herein.

 The present invention further provides a method of delivering a nucleic acid to a cell in a subject having neutralizing antibodies to AAV2 comprising administering to the subject an AAV5 particle containing a vector comprising the nucleic acid, thereby
25 delivering the nucleic acid to a cell in the subject. A subject that has neutralizing antibodies to AAV2 can readily be determined by any of several known means, such as contacting AAV2 protein(s) with an antibody-containing sample, such as blood, from a subject and detecting an antigen-antibody reaction in the sample. Delivery of the AAV5 particle can be by either *ex vivo* or *in vivo* administration as herein described.
30 Thus, a subject who might have an adverse immunogenic reaction to a vector

09717789-112100

administered in an AAV2 viral particle can have a desired nucleic acid delivered using an AAV5 particle. This delivery system can be particularly useful for subjects who have received therapy utilizing AAV2 particles in the past and have developed antibodies to AAV2. An AAV5 regimen can now be substituted to deliver the desired
5 nucleic acid.

In any of the methods of delivering heterologous nucleic acids to a cell or subject described herein, the AAV5-conjugated nucleic acid or AAV5 particle-conjugated nucleic acids described herein can be used.
10

In vivo administration to a human subject or an animal model can be by any of many standard means for administering viruses, depending upon the target organ, tissue or cell. Virus particles can be administered orally, parenterally (*e.g.*, intravenously), by intramuscular injection, by direct tissue or organ injection, by intraperitoneal injection,
15 topically, transdermally, via aerosol delivery, via the mucosa or the like. Viral nucleic acids (non-encapsidated) can also be administered, *e.g.*, as a complex with cationic liposomes, or encapsulated in anionic liposomes. The present compositions can include various amounts of the selected viral particle or non-encapsidated viral nucleic acid in combination with a pharmaceutically acceptable carrier and, in addition, if desired, may
20 include other medicinal agents, pharmaceutical agents, carriers, adjuvants, diluents, etc. Parental administration, if used, is generally characterized by injection. Injectables can be prepared in conventional forms, either as liquid solutions or suspensions, solid forms suitable for solution or suspension in liquid prior to injection, or as emulsions. Dosages will depend upon the mode of administration, the disease or condition to be treated, and
25 the individual subject's condition, but will be that dosage typical for and used in administration of other AAV vectors, such as AAV2 vectors. Often a single dose can be sufficient; however, the dose can be repeated if desirable.

Administration methods can be used to treat brain disorders such as Parkinson's
30 disease, Alzheimer's disease, and demyelination disease. Other diseases that can be

09717789-112100

treated by these methods include metabolic disorders such as , musculoskeletal diseases, cardiovascular disease, cancer, and autoimmune disorders.

Administration of this recombinant AAV5 virion to the cell can be
5 accomplished by any means, including simply contacting the particle, optionally contained in a desired liquid such as tissue culture medium, or a buffered saline solution, with the cells. The virion can be allowed to remain in contact with the cells for any desired length of time, and typically the virion is administered and allowed to remain indefinitely. For such *in vitro* methods, the virion can be administered to the
10 cell by standard viral transduction methods, as known in the art and as exemplified herein. Titers of virus to administer can vary, particularly depending upon the cell type, but will be typical of that used for AAV transduction in general which is well known in the art. Additionally the titers used to transduce the particular cells in the present examples can be utilized.

15

The cells that can be transduced by the present recombinant AAV5 virion can include any desired cell, such as the following cells and cells derived from the following tissues, human as well as other mammalian tissues, such as primate, horse, sheep, goat, pig, dog, rat, and mouse: Adipocytes, Adenocyte, Adrenal cortex, Amnion,
20 Aorta, Ascites, Astrocyte, Bladder, Bone, Bone marrow, Brain, Breast, Bronchus, Cardiac muscle, Cecum, Cervix, Chorion, Colon, Conjunctiva, Connective tissue, Cornea, Dermis, Duodenum, Endometrium, Endothelium, Endothelial cells, Epithelial tissue, Epithelial cells, Epidermis, Esophagus, Eye, Fascia, Fibroblasts, Foreskin, Gastric, Glial cells, Glioblast, Gonad, Hepatic cells, Histocyte, Ileum, Intestine, small
25 Intestine, Jejunum, Keratinocytes, Kidney, Larynx, Leukocytes, Lipocyte, Liver, Lung, Lymph node, Lymphoblast, Lymphocytes, Macrophages, Mammary alveolar nodule, Mammary gland, Mastocyte, Maxilla, Melanocytes, Mesenchymal, Monocytes, Mouth, Myelin, Myoblasts Nervous tissue, Neuroblast, Neurons, Neuroglia, Osteoblasts, Osteogenic cells, Ovary, Palate, Pancreas, Papilloma, Peritoneum, Pituicytes, Pharynx,
30 Placenta, Plasma cells, Pleura, Prostate, Rectum, Salivary gland, Skeletal muscle, Skin,

09717789.112100

Smooth muscle, Somatic, Spleen, Squamous, Stomach, Submandibular gland,
Submaxillary gland, Synoviocytes, Testis, Thymus, Thyroid, Trabeculae, Trachea,
Turbinate, Umbilical cord, Ureter, and Uterus.

5

STATEMENT OF UTILITY

09717789.112100

The present invention provides recombinant vectors based on AAV5. Such
vectors may be useful for transducing erythroid progenitor cells or cells lacking heparin
sulfate proteoglycans which is very inefficient with AAV2 based vectors. These
10 vectors may also be useful for transducing cells with a nucleic acid of interest in order
to produce cell lines that could be used to screen for agents that interact with the gene
product of the nucleic acid of interest. In addition to transduction of other cell types,
transduction of erythroid cells would be useful for the treatment of cancer and genetic
diseases which can be corrected by bone marrow transplants using matched donors.
15 Some examples of this type of treatment include, but are not limited to, the introduction
of a therapeutic gene such as genes encoding interferons, interleukins, tumor necrosis
factors, adenosine deaminase, cellular growth factors such as lymphokines, blood
coagulation factors such as factor VIII and IX, cholesterol metabolism uptake and
transport protein such as EpoE and LDL receptor, and antisense sequences to inhibit
20 viral replication of, for example, hepatitis or HIV.

The present invention provides a vector comprising the AAV5 virus as well as
AAV5 viral particles. While AAV5 is similar to AAV2, the two viruses are found
herein to be physically and genetically distinct. These differences endow AAV5 with
25 some unique advantages which better suit it as a vector for gene therapy.

Furthermore, as shown herein, AAV5 capsid protein is distinct from AAV2
capsid protein and exhibits different tissue tropism. AAV2 and AAV5 likely utilize
distinct cellular receptors. AAV2 and AAV5 are serologically distinct and thus, in a
30 gene therapy application, AAV5 would allow for transduction of a patient who already

possess neutralizing antibodies to AAV2 either as a result of natural immunological defense or from prior exposure to AAV2 vectors.

The present invention is more particularly described in the following examples
5 which are intended as illustrative only since numerous modifications and variations therein will be apparent to those skilled in the art.

EXAMPLES

10 To understand the nature of AAV5 virus and to determine its usefulness as a vector for gene transfer, it was cloned and sequenced.

Cell culture and virus propagation

Cos and HeLa cells were maintained as monolayer cultures in D10 medium
15 (Dulbecco's modified Eagle's medium containing 10% fetal calf serum, 100 µg/ml penicillin, 100 units/ml streptomycin and IX Fungizone as recommended by the manufacturer; (GIBCO, Gaithersburg, MD, USA) . All other cell types were grown under standard conditions which have been previously reported.

20 Virus was produced as previously described for AAV2 using the Beta galactosidase vector plasmid and a helper plasmid containing the AAV5 Rep and Cap genes (9). The helper plasmid was constructed in such a way to minimize any homologous sequence between the helper and vector plasmids. This step was taken to minimize the potential for wild-type (wt) particle formation by homologous
25 recombination.

DNA Cloning and Sequencing and Analysis

In order to clone the genome of AAV5, infectious cell lysate was expanded in adherent cos cells and then suspension HeLa cells with the resulting viral particles
30 isolated by CsCl isopycnic gradient centrifugation. DNA dot blots of Aliquots of the

09747789.112400

gradient fractions indicated that the highest concentration of viral genomes were contained in fractions with a refractive index of approx. 1.372. While the initial description of the virus did not determine the density of the particles, this value is similar to that of AAV2. Analysis of annealed virion derived DNA obtained from these

5 fractions indicated a major species of 4.6 kb in length which upon restriction analysis gave bands similar in size to those previously reported. Additional restriction mapping indicated a unique BssHII site at one end of the viral genome. This site was used to clone the major fragment of the viral genome. Additional overlapping clones were isolated and the sequence determined. Two distinct open reading frames (ORF) were

10 identified. Computer analysis indicated that the left-hand ORF is approx 60% similar to that of the Rep gene of AAV2. Of the 4 other reported AAV serotypes, all have greater than 90% similarity in this ORF. The right ORF of the viral capsid proteins is also approximately 60% homologous to the Capsid ORF of AAV2. As with other

15 AAV serotypes reported, the divergence between AAV5 and AAV2 is clustered in multiple blocks. By using the published three dimensional structure of the canine parvovirus and computer aided sequence comparisons, a number of these divergent regions have been shown to be on the exterior of the virus and thus suggest an altered tissue tropism.

20 Within the p5 promoter, a number of the core transcriptional elements are conserved such as the tataa box and YY1 site around the transcriptional start site. However the YY1 site at -60 and the upstream E-Box elements are not detectable suggesting an alternative method of regulation or activation.

25 The inverted terminal repeats (ITRs) of the virus were cloned as a fragment from the right end of the genome. The resulting fragment was found to contain a number of sequence changes compared to AAV2. However, these changes were found to be complementary and did not affect the ability of this region to fold into a hairpin structure. Within the stem region of the hairpin two sequence elements have been

30 found to be critical for the function of the ITRs as origins of viral replication. A repeat

00747789 "112100

motif of GAGC/T which serves as the recognition site of Rep and a GGTTGAG sequence downstream of the Rep binding site which is the position of Rep's site and strand specific cleavage reaction. This sequence is not conserved between AAV5 and the other cloned AAV's suggesting that the ITRs and Rep proteins of AAV5 cannot
5 compliment the other known AAV's.

To test the cross complementarity of AAV2 ITR containing genome and AAV5 ITR containing genomes recombinant particles were packaged either using type 2 Rep and Cap or type 5 Rep and Cap expression plasmids as previously described. As shown
10 in Fig. 2, viral particles were produced only when the respective expression plasmids were used to package the cognate ITRs. This result is distinct from that of other serotypes of AAV which have shown cross complementary in packaging.

This specificity of AAV5 Rep for AAV5 ITRs was confirmed using a terminal
15 resolution assay which can identify the site within one ITR cleaved by the Rep protein. Incubation of the Type 5 Rep protein with a type 2 ITR did not produce any cleavage products. In contrast, addition of type 2 Rep cleaved the DNA at the expected site. However AAV5 Rep did produce cleavage products when incubated with a type 5 ITR. The site mapped to a region 21 bases from the Rep binding motif that is similar to
20 AAV2 TRS. The site in AAV2 is CGGT TGAG (SEQ ID NO: 22) but in type 5 ITR is CGGT GTGA (SEQ ID NO: 21). The ability of AAV5 Rep to cleave at a different but similarly positioned site may result in integration of AAV5 at a distinct chromosomal locus compared to AAV2.

25 Recombinant virus produced using AAV5 Rep and Cap was obtained at a greater titer than type 2. For example, in a comparative study, virus was isolated from 8×10^7 COS cells by CsCl banding and the distribution of the Beta galactosidase genomes across the gradient were determined by DNA dot blots of aliquots of gradient fractions. DNA dot blot titers indicated that AAV5 particles were produced at a 10-50
30 fold higher level than AAV2.

09717789 112100

The sequence divergence in the capsid protein ORF implies that the tissue tropism of AAV2 and AAV5 would differ. To study the transduction efficiency of AAV5 and AAV2, a variety of cell lines were transduced with serial dilution's of the purified virus expressing the gene for nuclear localized Beta galactosidase activity.

5 Approx. 2×10^4 cells were exposed to virus in 1 ml of serum containing media for a period of 48-60 hrs. After this time the cells were fixed and stained for Beta-galactosidase activity with 5-Bromo-4-chloro- 3-indolyl-b-D- galactopyranoside (Xgal) (ICN Biochemicals). Biological titers were determined by counting the number of positive cells in the different dilutions using a calibrated microscope ocular then
10 multiplying by the area of the well. Titters were determined by the average number of cells in a minimum of 10 fields/well. Transduction of cos, HeLa, and 293, and IB3 cells with a similar number of particles showed approximately 10 fold decrease in titer with AAV5 compared with AAV2. In contrast MCF7 cells showed a 50-100 fold difference in transduction efficiency. Furthermore, both vectors transduced NIH 3T3
15 cells relatively poorly.

A recent publication reported that heparin proteoglycans on the surface of cells are involved in viral transduction. Addition of soluble heparin has been shown to inhibit transduction by blocking viral binding. Since the transduction data suggested a
20 difference in tissue tropism for AAV5 and AAV2, the sensitivity of AAV5 transduction to heparin was determined. At an MOI of 100, the addition of $20 \mu\text{g/ml}$ of heparin had no effect on AAV5 transduction. In contrast this amount of heparin inhibited 90% of the AAV2 transduction. Even at an MOI of 1000, no inhibition of AAV5 transduction was detected. These data support the conclusions of the tissue tropism study, i.e. that
25 AAV2 and AAV5 may utilize a distinct cell surface molecules and therefore the mechanism of uptake may differ as well.

AAV5 is a distinct virus within the dependovirus family based on sequence analysis, tissue tropism, and sensitivity to heparin. While elements of the P5 promoter
30 are retained between AAV2-6 some elements are absent in AAV5 suggesting

09747789-112100

alternative mechanism of regulation. The ITR and Rep ORF are distinct from those previously identified and fail to complement the packaging of AAV2 based genomes. The ITR of AAV5 contains a different TRS compared to other serotypes of AAV which is responsible for the lack of complementation of the ITRs. This unique TRS should also result in a different integration locus for AAV5 compared to that of AAV2. Furthermore the production of recombinant AAV5 particles using standard packaging systems is approx. 10-50 fold better than AAV2. The majority of the differences in the capsid proteins lies in regions which have been proposed to be on the exterior of the surface of the parvovirus. These changes are most likely responsible for the lack of cross reactive antibodies and altered tissue tropism compared to AAV2.

From the Rep ORF of AAV2, 4 proteins are produced; The p5 promoter (SEQ ID NO: 18) produces rep 68 (a spliced site mutant) and rep78 and the p19 promoter (SEQ ID NO: 16) produces rep 40 (a spliced site mutant) and rep 52. While these regions are not well conserved within the Rep ORF of AAV5 some splice acceptor and donor sites exist in approximately the same region as the AAV2 sites. These sites can be identified using standard computer analysis programs such as signal in the PCGENE program. Therefore the sequences of the Rep proteins can be routinely identified as in other AAV serotypes.

Hemagglutination assay

Hemagglutination activity was measured essentially as described previously (Chiorini et al 1997 J. Virol. Vol 71 6823-6833) Briefly 2 fold serial dilutions of virus in EDTA-buffered saline were mixed with an equal volume of 0.4% red blood cells in plastic U-bottom 96 well plates. The reaction was complete after a 2-h incubation at 8°C. Addition of purified AAV5 to a hemagglutination assay resulted in hemagglutination activity.

Transduction of airway epithelial cells

Primary airway epithelial cells were cultured and plated as previously described (Fasbender et al. J. Clin Invest. 1998 Jul 1; 102 (1): 184-93). Cells were transduced with an equivalent number of rAAV2 or rAAV5 particles containing a nuclear localized β -gal transgene with 50 particles of virus/cell (MOI 50) and continued in culture for 10 days. β -gal activity was determined following the procedure of (Chiorini et al. 1995 HGT Vol: 6 1531-1541) and the relative transduction efficiency compared. As shown in Figure 7, AAV5 transduced these cells 50- fold more efficiently than AAV2. This is the first time apical cells or cells exposed to the air have been shown to be infected by a gene therapy agent.

Transduction of striated muscle

Chicken myoblasts were cultured and plated as previously described (Rhodes & Yamada 1995 NAR Vol 23 (12) 2305-13). Cells were allowed to fuse and then transduced with a similar number of particles of rAAV2 or rAAV5 containing a nuclear localized β -gal transgene as previously described above after 5 days in culture. The cells were stained for β -gal activity following the procedure of (Chiorini et al. 1995 HGT Vol: 6 1531-1541) and the relative transduction efficiency compared. As shown in Figure 8, AAV5 transduced these cells approximately 16 fold more efficiently than AAV2.

Transduction of rat brain explants

Primary neonatal rat brain explants were prepared as previously described (Scortegagna et al. Neurotoxicology. 1997; 18 (2): 331-9). After 7 days in culture, cells were transduced with a similar number of particles of rAAV5 containing a nuclear localized β -gal transgene as previously described. After 5 days in culture, the cells were stained for β -gal activity following the procedure of (Chiorini et al. 1995 HGT Vol: 6 1531-1541). As shown in Figure 9, transduction was detected in a variety of cell types including astrocytes, neuronal cells and glial cells.

Transduction of human umbilical vein endothelial cells

Human umbilical vein endothelial cells were cultured and plated as previously described (Gnantenko et al. J Investig Med. 1997 Feb; 45(2): 87-98). Cells were transduced with rAAV2 or rAAV5 containing a nuclear localized β -gal transgene with 10 particles of virus/ cell (MOI 5) in minimal media then returned to complete media. After 24 hrs in culture the cells were stained for β -gal activity following the procedure of Chiorini et al. (1995 HGT Vol: 6 1531-1541), and the relative transduction efficiency compared. As shown in Figure 10, AAV5 transduced these cell 5-10 fold more efficiently than AAV2.

Throughout this application, various publications are referenced. The disclosures of these publications in their entireties are hereby incorporated by reference into this application in order to more fully describe the state of the art to which this invention pertains.

Although the present process has been described with reference to specific details of certain embodiments thereof, it is not intended that such details should be regarded as limitations upon the scope of the invention except as and to the extent that they are included in the accompanying claims.

References:

1. Arella, M., S. Garzon, J. Bergeron, and P. Tijssen. Handbook of Parvoviruses. Vol. 1. ed. P. Tijssen. Boca Raton, Florida, CRC Press, 1990.
2. Bachmann, P.A., M.D. Hoggan, E. Kurstak, J.L. Melnick, H.G. Pereira, P. Tattersall, and C. Vago. 1979. *Intervirology* 11: 248-254.
3. Bantel-Schaal, U. and M. Stohr. 1992. *J. Virol.* 66: 773-779.
4. Chang, L.S., Y. Shi, and T. Shenk. 1989. *J. Virol.* 63: 3479-88.
5. Chejanovsky, N. and B.J. Carter. 1989. *Virology* 173: 120-128.
6. Chejanovsky, N. and B.J. Carter. 1989. *Virology* 171: 239-247.

7. Chiorini, J.A., S.M. Wiener, R.M. Kotin, R.A. Owens, SRM Kyöstiö, and B. Safer. 1994. *J. Virol.* **68**: 7448-7457.
8. Chiorini, J.A., M.D. Weitzman, R.A. Owens, E. Urcelay, B. Safer, and R.M. Kotin. 1994. *J. Virol.* **68**: 797-804.
9. Chiorini, J.A., C.M. Wendtner, E. Urcelay, B. Safer, M. Hallek, and R.M. Kotin. 1995. *Human Gene Therapy* **6**: 1531-1541.
10. Chiorini, J.A., L. Yang, B. Safer, and R.M. Kotin. 1995. *J. Virol.* **69**: 7334-7338.
11. Dixit, M., M.S. Webb, W.C. Smart, and S. Ohi. 1991. *Gene* **104**: 253-7.
12. Fisher, R.E. and H.D. Mayor. 1991. *J Theor Biol* **149**: 429-39.
13. Flotte, T.R., S.A. Afione, C. Conrad, S.A. McGrath, R. Solow, H. Oka, P.L. Zeitlin, W.B. Guggino, and B.J. Carter. 1993. *Proc. Natl. Acad. Sci.* **90**: 10613-10617.
14. Flotte, T.R., S.A. Afione, R. Solow, M.L. Drumm, D. Markakis, W.B. Guggino, P.L. Zeitlin, and B.J. Carter. 1993. *J Biol Chem* **268**: 3781-90.
15. Hermonat, P.L., M.A. Labow, R. Wright, K.I. Berns, and N. Muzyczka. 1984. *J. Virol.* **51**: 329-339.
16. Hermonat, P.L. and N. Muzyczka. 1984. *Proc Natl Acad Sci USA* **81**: 6466-70.
17. Hunter, L.A. and R.J. Samulski. 1992. *J. Virol.* **66**: 317-24.
18. Ito, M. and H.D. Mayor. 1968. *J. Immuno.* **100**: 61-68.
19. Janik, J.E., M.M. Huston, K. Cho, and J.A. Rose. 1989. *Virology* **168**: 320-9.
20. Kaplitt, M.G., P. Leone, R.J. Samulski, X. Xiao, D.W. Pfaff, K.L. O'Malley, and J.M. During. 1994. *Nature Genetics* **8**: 148-154.
21. Kotin, R.M., M. Siniscalco, R.J. Samulski, X. Zhu, L. Hunter, C.A. Laughlin, S. McLaughlin, N. Muzyczka, M. Rocchi, and K.I. Berns. 1990. *Proc. Natl. Acad. Sci. (USA)* **87**: 2211-2215.
22. Laughlin, C.A., N. Jones, and B.J. Carter. 1982. *J. Virol.* **41**: 868-76.

09717789-112100

23. Laughlin, C.A., M.W. Myers, D.L. Risin, B.J. Carter. 1979. *Virology* **94**: 162-74.
24. McCarty, D.M., J. Pereira, I. Zolotukhin, X. Zhou, J.H. Ryan, and N. Muzyczka. 1994. *J. Virol.* **68**: 4988-4997.
25. Mendelson, E., J.P. Trempe, and B.J. Carter. 1986. *J. Virol.* **60**: 823-832.
26. Mizukami, H., N.S. Young, and K.E. Brown. 1996. *Virology* **217**: 124-130.
27. Muster, C.J., Y.S. Lee, J.E. Newbold, and J. Leis. 1980. *J. Virol.* **35**: 653-61.
28. Muzyczka, N. 1992. *Curr Top Microbiol Immunol* **158**: 97-129.
29. Parks, W.P., J.L. Melnick, R. Rongey, and H.D. Mayor. 1967. *J. Virol.* **1**: 171-180.
30. Podsakoff, G., K.K. Jr Wong, and S. Chatterjee. 1994. *J. Virol.* **68**: 5656-5666.
31. Rose, J.A., M.D. Hoggan, F. Koczot, and A.J. Shatkin. 1968. *J. Virol.* **2**: 999-1005.
32. Russell, D.W., A.D. Miller, and I.E. Alexander. 1994. *Proc. Natl. Acad. Sci. USA* **91**: 8915-8919.
33. Ryan, J.H., S. Zolotukhin, and N. Muzyczka. 1996. *J. Virol.* **70**: 1542-1553.
34. Samulski, R.J., K.I. Berns, M. Tan, and N. Muzyczka. 1982. *Proc Natl Acad Sci USA* **79**: 2077-81.
35. Samulski, R.J., L.S. Chang, and T. Shenk. 1989. *J. Virol.* **63**: 3822-8.
36. Sanes, J.R., J.L.R. Rubenstein, and J.F. Nicocas. 1986. *EMBO* **5**: 3133-3142.
37. Senapathy, P., J.D. Tratschin, and B.J. Carter. 1984. *J Mol Biol* **179**: 1-20.
38. Tratschin, J.D., I.L. Miller, and B.J. Carter. 1984. *J. Virol.* **51**: 611-619.
39. Trempe, J.P. and B.J. Carter. 1988. *J. Virol.* **62**: 68-74.
40. Trempe, J.P., E. Mendelson, and B.J. Carter. 1987. *Virology* **161**: 18-28.
41. Walsh, C.E., J.M. Liu, X. Xiao, N.S. Young, A.W. Nienhuis, and R.J. Samulski. 1992. *Proc Natl Acad Sci USA* **89**: 7257-61.

09717789 112100

42. **Winocour, E., M.F. Callaham, and E. Huberman.** 1988. *Virology* **167**: 393-9.
43. **Jaksch, M., K.D. Gerbitz, and C. Kilger.** 1995. *Clin. Biochem.* **28**:503-509
44. **Burcin, M.M., O'Malley, B.W. and S.Y. Tsai.** 1998. *Frontiers in Bioscience* **3**:1-7.

001211" 68221260